

## Automata and Formal Languages — Exercise Sheet 13

### Exercise 13.1

Let  $AP = \{p, q\}$  and let  $\Sigma = 2^{AP}$ . Give LTL formulas for the following  $\omega$ -languages:

- (a)  $\{p, q\} \emptyset \Sigma^\omega$
- (b)  $\Sigma^* \{q\}^\omega$
- (c)  $\Sigma^* (\{p\} + \{p, q\}) \Sigma^* \{q\} \Sigma^\omega$
- (d)  $\{p\}^* \{q\}^* \emptyset^\omega$

In (a) and (d) the  $\emptyset$  symbol stands for the letter  $\emptyset \in \Sigma$ , and not for the empty  $\omega$ -language.

### Exercise 13.2

Let  $AP = \{p, q\}$  and let  $\Sigma = 2^{AP}$ . Give Büchi automata for the  $\omega$ -languages over  $\Sigma$  defined by the following LTL formulas:

- (a)  $\mathbf{XG}\neg p$
- (b)  $(\mathbf{GF}p) \rightarrow (\mathbf{F}q)$
- (c)  $p \wedge \neg(\mathbf{XF}p)$
- (d)  $\mathbf{G}(p \mathbf{U} (p \rightarrow q))$
- (e)  $\mathbf{F}q \rightarrow (\neg q \mathbf{U} (\neg q \wedge p))$

### Exercise 13.3

Say which of the following equivalences hold. For every equivalence that does not hold give an instantiation of  $\varphi$  and  $\psi$  together with a computation that disproves the equivalence.

- (a)  $\mathbf{F}(\varphi \vee \psi) \equiv \mathbf{F}\varphi \vee \mathbf{F}\psi$
- (b)  $\mathbf{F}(\varphi \wedge \psi) \equiv \mathbf{F}\varphi \wedge \mathbf{F}\psi$
- (c)  $\mathbf{G}(\varphi \vee \psi) \equiv \mathbf{G}\varphi \vee \mathbf{G}\psi$
- (d)  $(\varphi \vee \psi) \mathbf{U} \rho \equiv (\varphi \mathbf{U} \rho) \vee (\psi \mathbf{U} \rho)$
- (e)  $\mathbf{GF}(\varphi \wedge \psi) \equiv \mathbf{GF}\varphi \wedge \mathbf{GF}\psi$
- (f)  $\mathbf{X}(\varphi \mathbf{U} \psi) \equiv (\mathbf{X}\varphi \mathbf{U} \mathbf{X}\psi)$

### Exercise 13.4

Let  $AP = \{p, q\}$  and let  $\Sigma = 2^{AP}$ . An LTL formula is a tautology if it is satisfied by all computations. Which of the following LTL formulas are tautologies?


- (a)  $\mathbf{G}p \rightarrow \mathbf{F}p$
- (b)  $\mathbf{G}(p \rightarrow q) \rightarrow (\mathbf{G}p \rightarrow \mathbf{G}q)$
- (c)  $\mathbf{F}\mathbf{G}p \vee \mathbf{F}\mathbf{G}\neg p$
- (d)  $\neg\mathbf{F}p \rightarrow \mathbf{F}\neg\mathbf{F}p$
- (e)  $(\mathbf{G}p \rightarrow \mathbf{F}q) \leftrightarrow (p \mathbf{U} (\neg p \vee q))$
- (f)  $\neg(p \mathbf{U} q) \leftrightarrow (\neg p \mathbf{U} \neg q)$
- (g)  $\mathbf{G}(p \rightarrow \mathbf{X}p) \rightarrow (p \rightarrow \mathbf{G}p)$

**Solution 13.1**


- (a)  $(p \wedge q) \wedge \mathbf{X}(\neg p \wedge \neg q)$
- (b)  $\mathbf{FG}(\neg p \wedge q)$
- (c)  $\mathbf{F}(p \wedge \mathbf{XF}(\neg p \wedge q))$
- (d)  $(p \wedge \neg q) \mathbf{U} ((\neg p \wedge q) \mathbf{U} \mathbf{G}(\neg p \wedge \neg q))$

**Solution 13.2**


(a)




(b) Note that  $(\mathbf{GF}p) \rightarrow (\mathbf{F}q) \equiv \neg(\mathbf{GF}p) \vee (\mathbf{F}q) \equiv (\mathbf{FG}\neg p) \vee (\mathbf{F}q)$ . We construct Büchi automata for  $\mathbf{FG}\neg p$  and  $\mathbf{F}q$ , and take their union:




(c) Note that  $p \wedge \neg(\mathbf{XF}p) \equiv p \wedge \mathbf{XG}\neg p$ . We construct a Büchi automaton for  $p \wedge \mathbf{XG}\neg p$ :



(d)



(e) Note that  $\mathbf{F}q \rightarrow (\neg q \mathbf{U} (\neg q \wedge p)) \equiv \mathbf{G}\neg q \vee (\neg q \mathbf{U} (\neg q \wedge p))$ . Consider this case split over the occurrence of a  $p$ : computations that satisfy the formula either have no occurrence of  $p$ , in which case they must satisfy the first part of the  $\vee$  (i.e.  $\mathbf{G}\neg q$ ), or they have a first occurrence of  $p$  with no  $q$  before or at the same time:



**Solution 13.3**

(a) True, since:

$$\begin{aligned}
 \sigma \models \mathbf{F}(\varphi \vee \psi) &\iff \exists k \geq 0 \text{ s.t. } \sigma^k \models (\varphi \vee \psi) \\
 &\iff \exists k \geq 0 \text{ s.t. } (\sigma^k \models \varphi) \vee (\sigma^k \models \psi) \\
 &\iff (\exists k \geq 0 \text{ s.t. } \sigma^k \models \varphi) \vee (\exists k \geq 0 \text{ s.t. } \sigma^k \models \psi) \\
 &\iff \sigma \models \mathbf{F}\varphi \vee \mathbf{F}\psi.
 \end{aligned}$$

(b) False. Let  $\sigma = \{p\}\{q\}\emptyset^\omega$ . We have  $\sigma \models \mathbf{F}p \wedge \mathbf{F}q$  and  $\sigma \not\models \mathbf{F}(\varphi \wedge \psi)$ .

(c) False. Let  $\sigma = (\{p\}\{q\})^\omega$ . We have  $\sigma \models \mathbf{G}(p \vee q)$  and  $\sigma \not\models \mathbf{G}p \vee \mathbf{G}q$ .

(d) False. Let  $\sigma = \{p\}\{q\}\{r\}\emptyset^\omega$ . We have  $\sigma \models (p \vee q) \mathbf{U} r$  and  $\sigma \not\models (p \mathbf{U} r) \vee (q \mathbf{U} r)$ .

(e) False. Let  $\sigma = (\{p\}\{q\})^\omega$ . We have  $\sigma \not\models \mathbf{GF}(p \wedge q)$  and  $\sigma \models \mathbf{GF}p \wedge \mathbf{GF}q$ .

(f) True, since:

$$\begin{aligned}
 \sigma \models \mathbf{X}(\varphi \mathbf{U} \psi) &\iff \sigma^1 \models (\varphi \mathbf{U} \psi) \\
 &\iff \exists k \geq 0 : (\sigma^1)^k \models \varphi \text{ and } \forall 0 \leq i < k (\sigma^1)^i \models \psi \\
 &\iff \exists k \geq 0 : (\sigma^k)^1 \models \varphi \text{ and } \forall 0 \leq i < k (\sigma^i)^1 \models \psi \\
 &\iff \exists k \geq 0 : \sigma^k \models \mathbf{X}\varphi \text{ and } \forall 0 \leq i < k (\sigma^i \models \mathbf{X}\psi) \\
 &\iff \sigma \models (\mathbf{X}\varphi) \mathbf{U} (\mathbf{X}\psi).
 \end{aligned}$$

**Solution 13.4**

(a)  $\mathbf{G}p \rightarrow \mathbf{F}p$  is a tautology since

$$\begin{aligned}
 \sigma \models \mathbf{G}p &\iff \forall k \geq 0 \sigma^k \models p \\
 &\implies \exists k \geq 0 \sigma^k \models p \\
 &\iff \sigma \models \mathbf{F}p.
 \end{aligned}$$

(b)  $\mathbf{G}(p \rightarrow q) \rightarrow (\mathbf{G}p \rightarrow \mathbf{G}q)$  is a tautology. For the sake of contradiction, suppose this is not the case. There exists  $\sigma$  such that

$$\sigma \models \mathbf{G}(p \rightarrow q), \text{ and} \tag{1}$$

$$\sigma \not\models (\mathbf{G}p \rightarrow \mathbf{G}q). \tag{2}$$

By (2), we have

$$\begin{aligned}
 \sigma \models \mathbf{G}p, \text{ and} \\
 \sigma \not\models \mathbf{G}q.
 \end{aligned}$$

Therefore, there exists  $k \geq 0$  such that  $p \in \sigma(k)$  and  $q \notin \sigma(k)$  which contradicts (1).

(c)  $\mathbf{F}\mathbf{G}p \vee \mathbf{F}\mathbf{G}\neg p$  is not a tautology since it is not satisfied by  $(\{p\}\{q\})^\omega$ .

(d)  $\neg\mathbf{F}p \rightarrow \mathbf{F}\neg\mathbf{F}p$  is a tautology since  $\varphi \rightarrow \mathbf{F}\varphi$  is a tautology for every formula  $\varphi$ .

(e)  $(\mathbf{G}p \rightarrow \mathbf{F}q) \leftrightarrow (p \mathbf{U} (\neg p \vee q))$  is a tautology. We have

$$\begin{aligned}
 \mathbf{G}p \rightarrow \mathbf{F}q &\equiv \neg\mathbf{G}p \vee \mathbf{F}q && \text{(by def. of implication)} \\
 &\equiv \mathbf{F}\neg p \vee \mathbf{F}q \\
 &\equiv \mathbf{F}(\neg p \vee q) \\
 &\equiv \mathbf{F}(p \rightarrow q) && \text{(by def. of implication)}
 \end{aligned}$$

Therefore, we have to show that

$$\mathbf{F}(p \rightarrow q) \leftrightarrow (p \mathbf{U} (p \rightarrow q)).$$

$\leftarrow$ ) Let  $\sigma$  be such that  $\sigma \models (p \mathbf{U} (p \rightarrow q))$ . In particular, there exists  $k \geq 0$  such that  $\sigma^k \models (p \rightarrow q)$ . Therefore,  $\sigma \models \mathbf{F}(p \rightarrow q)$ .

$\rightarrow$ ) Let  $\sigma$  be such that  $\sigma \models \mathbf{F}(p \rightarrow q)$ . Let  $k \geq 0$  be the smallest position such that  $\sigma^k \models (p \rightarrow q)$ . For every  $0 \leq i < k$ , we have  $\sigma^i \not\models (p \rightarrow q)$  which is equivalent to  $\sigma^i \models p \wedge \neg q$ . Therefore, for every  $0 \leq i < k$ , we have  $\sigma^i \models p$ . This implies that  $\sigma \models p \mathbf{U} (p \rightarrow q)$ .

(f)  $\neg(p \mathbf{U} q) \leftrightarrow (\neg p \mathbf{U} \neg q)$  is not a tautology. Let  $\sigma = \{p\}\{q\}^\omega$ . We have  $\sigma \not\models \neg(p \mathbf{U} q)$  and  $\sigma \models (\neg p \mathbf{U} \neg q)$ .

(g)  $\mathbf{G}(p \rightarrow \mathbf{X}p) \rightarrow (p \rightarrow \mathbf{G}p)$  is a tautology since

$$\begin{aligned}
 \mathbf{G}(p \rightarrow \mathbf{X}p) \rightarrow (p \rightarrow \mathbf{G}p) &\equiv \neg \mathbf{G}(\neg p \vee \mathbf{X}p) \vee (\neg p \vee \mathbf{G}p) && \text{(by def. of implication)} \\
 &\equiv \mathbf{F}(p \wedge \neg \mathbf{X}p) \vee \neg p \vee \mathbf{G}p \\
 &\equiv \neg \mathbf{G}p \rightarrow (\neg p \vee (\mathbf{F}(p \wedge \mathbf{X}\neg p))) && \text{(by def. of implication)} \\
 &\equiv \mathbf{F}\neg p \rightarrow (\neg p \vee (\mathbf{F}(p \wedge \mathbf{X}\neg p))) \\
 &\equiv \mathbf{F}\neg p \rightarrow \mathbf{F}\neg p.
 \end{aligned}$$